

Age and Gender-Related Variations in Third Ventricle Morphology: An MRI Analysis

Dr. V. Sai Abhinay

Post graduate, Department of Radio diagnosis, ACS Medical college and hospital, Chennai

Dr. Saranya Ayyadurai

Assistant Professor Department of Radio diagnosis, ACS Medical college and hospital, Chennai

***Dr. Vinoth Thangam**

Associate Professor Department of Radio diagnosis, ACS Medical college and hospital, Chennai

Abstract:

Background: The third ventricle, critical for cerebrospinal fluid dynamics and located near crucial neuroendocrine regions like the hypothalamus and thalamus, shows variations in morphology influenced by age and gender. These variations may relate to neurological conditions and cognitive changes, necessitating a deeper understanding through Magnetic Resonance Imaging (MRI).

Methods: This cross-sectional study involved 100 participants, stratified equally by gender and distributed among five age groups (20-70 years), without neurological or psychiatric disorders. High-resolution MRIs were used to assess the third ventricle dimensions. Width, height, and length measurements were analyzed using statistical methods to explore relationships with age and gender.

Results: The study found a moderate positive correlation between age and third ventricle width ($r = 0.45$, $p < 0.01$), indicating an increase in size with age. Additionally, males exhibited a slightly larger third ventricle width compared to females. These findings suggest that ventricular size may be a potential biomarker for aging and neurodegenerative processes, with gender differences possibly influencing disease susceptibility and progression.

Conclusion: The morphological changes in the third ventricle related to age and gender have significant implications for diagnosing and understanding the pathophysiology of neurodegenerative diseases. While insightful, the findings are limited by the study's sample size and its cross-sectional nature. Future research should expand on these results with a larger, more diverse cohort and consider longitudinal studies to better understand the implications of third ventricle morphology changes over time. The human brain, a complex and dynamic organ, undergoes various structural and functional changes across the lifespan, influenced by numerous factors including genetics, environment, health status, age, and gender. Among the structures of interest within the brain, the third ventricle, a narrow cavity located deep within the cerebral hemispheres, serves as a critical component of the brain's ventricular system, facilitating the flow of cerebrospinal fluid (CSF) throughout the brain and spinal cord. This ventricle's proximity to vital structures such as the hypothalamus and thalamus underscores its

importance in neuroendocrine regulation, making its study via Magnetic Resonance Imaging (MRI) of paramount importance for understanding a wide array of neurological conditions and the aging process itself [1].

Recent advancements in MRI technology have propelled forward our understanding of brain anatomy, allowing for the in-depth study of the third ventricle in unparalleled detail. The precise imaging capabilities of MRI not only facilitate the visualization of changes in ventricular size and shape but also enable the assessment of these changes in relation to age and gender [2]. This is particularly relevant given the growing body of evidence suggesting that the morphological characteristics of the third ventricle can vary significantly across individuals, potentially reflecting or even predicting cognitive changes, susceptibility to neurological diseases, and responses to therapies.

The correlation of the third ventricle's morphology with age and gender is not merely an academic curiosity but a crucial area of investigation with profound clinical implications. Age-related changes in ventricular size, for example, have been associated with cognitive decline and the development of neurodegenerative diseases such as Alzheimer's. Understanding these changes and their normative ranges is essential for distinguishing between healthy aging and early signs of pathology. Moreover, gender differences in brain structures have been observed, with potential implications for disease susceptibility, progression, and response to treatment. This underscores the necessity for gender-specific analysis in neurological research and clinical practice [3].

Building upon these considerations, this original article embarks on a comprehensive examination of the third ventricle's morphological characteristics as depicted by MRI, focusing on how these characteristics correlate with the individual's age and gender. By synthesizing findings from recent research, employing robust methodological approaches, and delving into the biological and clinical significance of observed patterns, this article aims to enrich our understanding of the third ventricle in the context of brain health and disease [4].

In pursuit of this aim, we will first review the existing literature to outline the current knowledge and gaps regarding the third ventricle's morphology and its implications. Following this, we will detail the methodologies employed in contemporary research for assessing the third ventricle via MRI, highlighting the techniques and criteria used for measurement and analysis. Subsequent sections will present and discuss the findings from our investigation, emphasizing the significance of age and gender differences in the third ventricle's morphology. Finally, we will explore the potential clinical applications of our findings, considering how they could inform diagnostic criteria, therapeutic strategies, and future research directions.

Through this detailed exploration, we aspire to contribute valuable insights to the fields of neurology and neuroscience, highlighting the critical role of individual differences in brain structure when assessing and addressing neurological health and disease.

Methodology

Setting

This study was conducted in the Department of Radiology at a tertiary care hospital. The department is equipped with advanced Magnetic Resonance Imaging (MRI) facilities, including a high-resolution 3T MRI scanner, which was utilized for obtaining detailed images of the brain, specifically focusing on the third ventricle.

Participants

A total of 100 participants were enrolled in this study through a convenience sampling method. The participants were divided equally among genders (50 males and 50 females) and stratified into five age groups (20-30, 31-40, 41-50, 51-60, 61-70 years), with 10 males and 10 females in each age group.

Inclusion Criteria

Participants aged between 20 and 70 years.

Participants willing to give informed consent for participation in the study.

Participants with no history of neurological or psychiatric disorders.

Participants without any contraindications to MRI, such as pacemakers, cochlear implants, or claustrophobia.

Exclusion Criteria

Participants with a history of neurosurgery or brain injury.

Participants diagnosed with any neurodegenerative diseases, such as Alzheimer's disease, Parkinson's disease, or multiple sclerosis.

Participants with a history of significant alcohol or substance abuse.

Participants with any known congenital abnormalities of the brain.

Participants with artifacts on MRI images due to movement or metal implants.

Procedure

MRI Acquisition

All participants underwent MRI scans using a 3T MRI scanner equipped with an 8-channel head coil. The imaging protocol was standardized to include high-resolution T1-weighted images with the following parameters: repetition time (TR) = 1900 ms, echo time (TE) = 2.26 ms, flip angle = 9°, field of view (FOV) = 256x256 mm, matrix size = 256x256, slice thickness = 1 mm, and no gap between slices. Special attention was given to obtaining high-quality sagittal and axial images to visualize the third ventricle clearly.

Image Analysis

The MRI images were analyzed by two experienced radiologists who were blinded to the participants' age and gender. The third ventricle's width, length, and height were measured using the hospital's standard DICOM viewer software. The width was measured at the widest part of the third ventricle on axial images, the height was measured from the floor to the roof in the sagittal plane, and the length was measured from the anterior to the posterior commissure in the sagittal plane. Any discrepancies between the measurements of the two radiologists were resolved through consensus.

Statistical Analysis

The data were analyzed using statistical software (SPSS version 26.0). Descriptive statistics were used to summarize the demographic characteristics of the study population and the


morphometric measurements of the third ventricle. The correlations between the third ventricle's dimensions and age, as well as gender differences, were assessed using Pearson's correlation coefficient and independent samples t-tests, respectively. A p-value of less than 0.05 was considered statistically significant.

Ethics Approval

This study was conducted in accordance with the Declaration of Helsinki and approved by the Institutional Review Board (IRB) of the hospital. Written informed consent was obtained from all participants before enrollment in the study.

Results

Demographic Characteristic	Total (N=100)	Male (n=50)	Female (n=50)
Age (years)			
20-30	20	10	10
31-40	20	10	10
41-50	20	10	10
51-60	20	10	10
61-70	20	10	10
Mean Age	45 ± 15	45 ± 15	45 ± 15
History of Neurological Disorders	0	0	0



This scatterplot shows the relationship between the participants' age and the width of the third ventricle. The x-axis represents the age of the participants, while the y-axis represents the width of the third ventricle in millimeters. Each point represents a participant, with male participants shown in blue and female participants in pink.

$r = 0.45, p < 0.01$, indicating a moderate positive correlation, suggesting that the third ventricle width tends to increase with age.




Figure 2: Shows the zoomed in and out views of the third ventricle

Discussion:

The findings of this study offer significant insights into the correlation between the third ventricle's dimensions and age and gender. Consistent with prior research, our results revealed a moderate positive correlation between age and the width of the third ventricle ($r = 0.45, p < 0.01$), suggesting that as individuals age, there is a discernible increase in the width of the third ventricle. This correlation may reflect the natural process of brain aging, including the potential loss of brain volume and changes in cerebrospinal fluid dynamics[5-7].

Interestingly, our study also found gender differences in the third ventricle's width, with males showing a slightly larger width on average than females ($t = -2.58, p < 0.05$). This finding

aligns with existing literature suggesting gender-specific differences in brain structures, which may have implications for understanding gender-specific risks for neurological diseases[6] .

The increase in third ventricle width with age observed in our study echoes findings from previous research, indicating that ventricular enlargement may serve as a biomarker for aging or neurodegenerative processes[8]. The precise mechanisms underlying these changes remain a subject of ongoing investigation, with potential factors including the differential impact of aging on brain parenchyma, variations in CSF dynamics, and the resilience of brain structures to age-related atrophy.

Furthermore, the gender differences observed in third ventricle dimensions in our cohort underline the importance of considering gender as a factor in neurological assessments and research. These differences might be attributable to variations in hormonal influences, genetic factors, or differential susceptibility to environmental factors, each of which could influence brain morphology [9,10].

Conclusion

Our study's findings should be interpreted in the context of its limitations, including the relatively small sample size and the cross-sectional design, which precludes causal inferences. Additionally, the use of a single MRI modality and measurement technique may limit the generalizability of our findings. Future research should consider longitudinal designs, larger and more diverse populations, and multimodal imaging approaches to further elucidate the relationship between third ventricle dimensions and neurological health.

References:

1. Patnaik P, Singh V, Singh D, Singh S. Gender Related Differences in Third Ventricle Parameters with Correlation to Cerebrum Size - A Study on Head CT Scans. *International Journal of Health Sciences and Research*. 2015;5:140-7.
2. Salk I, Atalar M, Sezer F, Eğılmez H, Çetin A, Arslan M. An MRI study of age-related changes in the dimensions related temporal lobe. *International journal of clinical and experimental medicine*. 2014;73:515-22.
3. Patnaik P, Singh S, Singh D, Singh V. Morphometric Study of the Third Ventricles in Apparently Normal Subjects Using Computerized Tomography. *International Journal of Health Sciences and Research*. 2016;6:153-9.
4. Singh S, Sharma B, Prajapati U, Sharma P, Bhatta M, Paudel N. Estimation of ventricles size of human brain by Magnetic Resonance Imaging in Nepalese Population: A retrospective study. 2020;13:45-50. DOI: 10.3126/jgmcn.v13i1.28542.
5. Khanal U, Sah K, Chand R. Assessment Of Third Ventricular Parameters in Computed Tomography of Head: A Pilot Study in Tertiary Level Hospital. *Journal of Institute of Medicine*. 2020;42:82-6. DOI: 10.3126/JIOM.V42I3.37594.
6. Ahmed S, Akram MN, Ashraf A, Noor A, Irshad F, Nishat M. CT Scan Study of Third Ventricles Dimensions in Normal Population. *Pakistan Journal of Medical and Health Sciences*. 2022. DOI: 10.53350/pjmhs20221611345.

7. Zhuravlova I, Kornieieva M. Anatomic Variability of the Morphometric Parameters of the Third Ventricle of the Brain and Its Relations to the Shape of the Skull. *Journal of Neurological Surgery Part B: Skull Base*. 2020;82:e315 - e320. DOI: 10.1055/s-0040-1701527.
8. Nazir M, Shah B, Aslam B, Amin R, Bhat S. Third Ventricle of Brain: A Morphometric Study by Computerized Tomography. *Paripex Indian Journal Of Research*. 2020.
9. Polat S, Öksüzler FY, Öksüzler M, Kabakci AG, Yücel A. Morphometric MRI Study of the Brain Ventricles in Healthy Turkish Subjects. *International Journal of Morphology*. 2019. DOI: 10.4067/S0717-95022019000200554.
10. Usman J, Zagga A, Tadros AA, Yunusa G, Saidu S, Ahmed H, Bello S, Abdulhameed A, Bello A. Morphological variation of third ventricle using computerized tomography among different gender and age groups: A 5-year retrospective study in Usmanu Danfodiyo University Teaching Hospital, Sokoto, North - West Nigeria. *Sahel Medical Journal*. 2013;16:83. DOI: 10.4103/1118-8561.121904.